

Bowel trouble *could be* all in the mind



NG WAN CHING

Experts are linking irritable bowel syndrome with psychological disorders but needing help for mental issues is often perceived as a stigma by patients

A 66-year-old housewife suffers from irritable bowel syndrome (IBS) as well as anxiety and a few other psychological issues.

With the help of a gastroenterologist and psychologist, the woman, who wanted to be known only as Madam Theresa, traced the cause of her psychological issues to when she suffered abuse as a child at a boarding school in another country.

She had not expected to discover that her bowel condition was probably caused by her psychological issues.

She had been suffering from IBS for years before a doctor suggested that a psychologist might be able to help.

Her physical symptoms include an intense pain between her ribs, bloatedness and gas. As a result, she eats very little.

She also suffers from low moods which have affected her relationship with her husband.

Recently, over the past year or so, she has been seeing a psychologist at National University Hospital (NUH) for her psychological symptoms.

Increasingly, doctors are associating psychological disorders with people who have IBS.

To detect mental issues early, doctors at Singapore General Hospital (SGH) have started to screen IBS patients for psychological disorders, while at NUH, a gastroenterologist and a psychologist both see IBS patients together.

IBS is one of the most common conditions that patients seek help for at NUH and SGH specialist clinics for gastrointestinal disorders.

A local study published about 10 years ago estimates the population prevalence at about 8 to 10 per cent, said Dr Reuben Wong, a consultant at the department of gastroenterology and hepatology at NUH.

Doctors are now working on making IBS a notifiable condition and having a registry on it, he said.

Madam Theresa's case is not atypical, he said. The early studies published by his mentor, Dr Douglas Drossman, from the University of North Carolina at Chapel Hill School of Medicine in the United States, describe a history of childhood abuse and issues in IBS patients.

"No one believed him years ago when he first presented his results, but it is now well established that there is a significantly higher history of abuse, neglect and trauma in IBS patients compared with healthy individuals," said Dr Wong.

After studies overseas proved that there is a link between IBS and psychological disorders, Singapore has done its own study.

Doctors at SGH found that almost half of 345 IBS patients screened from November 2010 to October 2011 also had psychological disorders.

In comparison, the 2010 Singapore Mental Health Study found that 12 per cent of people here will be stricken by mental illness in their lifetime.

The IBS patients found to have psychological disorders in the SGH study were then offered psychological consultations.

However, only half of them opted for them. Of these, half recovered from their psychological disorders and their quality of life improved.

This shows the importance of psychological intervention in the management of IBS patients with psychological disorders, said Dr Wang Yu Tien, a consultant at the department of gastroenterology and hepatology at SGH. He is the principal investigator of the study, which was submitted for SGH's Annual Scientific Meeting this month.

COMMONLY AFFECTS YOUNGER ADULTS

Although IBS is not life-threatening, it can negatively impact a patient's lifestyle and quality of life as well as that of his partner's.

Unlike diabetes or hypertension, it is not a disease which becomes more common with age. Instead, it commonly affects younger adults in the prime of their life – the most productive members in society – Dr Wong added.

There is also much overlap between IBS and other functional gastrointestinal disorders, so the diagnosis is not always clear cut.

The other challenge now, as the SGH study shows, is to get more IBS patients who have psychological issues to seek psychological help.

Much of the resistance such patients put up against seeking help is the perceived stigma of having a psychological or psychiatric disorder.

Dr Victor Kwok, an associate consultant at the department of psychiatry at SGH, said some IBS patients tell doctors that they are "not mad" when psychological help is suggested.

"People still think it is a weakness, character failing or within a person's control. But it is a medical condition just like diabetes or high blood pressure," he said.

Many patients also do not accept that their bowel symptoms have anything to do with their brain, said Dr Wang.

"They tell us it's two completely different things,



PHOTO ILLUSTRATION: LIM KAILI, ISTOCKPHOTO

two different organs. This stems from a lack of knowledge," he said.

But researchers are finding evidence that dysfunction along pathways running between the brain and the intestines may be contributing to the symptoms of IBS, which include abdominal pain, constipation and diarrhoea.

Nerves in the intestines that are experiencing excessive sensitivity can trigger changes in the brain.

Thoughts, feelings and activation of parts of the brain that have to do with anxiety or arousal can stimulate exaggerated responses from the gut.

What is not known is which comes first, said Dr Wang.

"It could be that the gut is not feeling well, then the brain feels depressed. Or the person is depressed, which then affects the gut," he said.

The more common psychiatric conditions that Dr Kwok has diagnosed in IBS patients are depression, dysthymia (low mood that has persisted for at least two years), generalised anxiety disorder, panic disorders and somatisation disorders in which the patient experiences multiple physical symptoms that are not explained by disease and primary insomnia.

Patients with psychological problems can develop counter-productive or unsuitable behaviour to do with eating or passing motion which may make the IBS worse, said Dr Kwok.

Others may tend to seek medical help more often, which can also worsen their condition.

"The more anxious patients may ask for more tests and need a lot more reassurance from doctors. This can aggravate the IBS and reduce their quality of life, said Dr Kwok.

Once it is established that the patient has psychological symptoms, he should get psychological help.

The belief is that reduction of psychological symptoms leads to the reduction of bowel symptoms.

Such treatment for IBS has good scientific evidence and is endorsed by many institutes including the American College of Gastroenterology, said Dr Kwok.

PSYCHOLOGICAL SCREENING

Since 2009, SGH has started an early intervention programme for anxiety and depression among IBS patients.

Since then, all IBS patients have the option of getting a free screening to detect depression and anxiety that might otherwise have gone undetected and untreated.

The psychological screening is done by an IBS coordinator, who goes through a few questionnaires with each IBS patient.

If a patient is suspected of having anxiety disorders or depression, he is given the option of seeing a psychiatrist or psychologist, and attending a support group for IBS patients, said Dr Kwok.

The support group, which holds meetings once every month at SGH, helps especially those who are afraid of the stigma of seeing a psychiatrist, he said.

Underscoring the importance of psychological help for patients, psychiatrists and psychologists at SGH will soon be working in much closer proximity to the gastroenterologists, in the same clinic.

An integrated multi-disciplinary IBS clinic will be run from next month, said Dr Wang.

This is already being done at NUH. In the Gastrointestinal-Psychology Collaborative Service, started in 2009, a psychologist sits in a clinic next to the gastroenterologist in case any IBS patient with psychological disorders wants a consultation.

Dr Wong started this together with the

department of psychiatry at NUH, in a collaboration with psychologists from James-Cook University Singapore.

At the time, he had just returned from his fellowship at the University of North Carolina Centre for Functional GI Disorders & Motility Disorders.

There, he found that there were dedicated psychologists who were on-site at the clinic, seeing the patients together with the doctors.

"This team-based approach allowed patients to be treated holistically by a gastroenterologist and psychologist working together," he said.

HOLISTIC TREATMENT APPROACH

With psychiatrists and psychologists on board, doctors are also trying to change the way IBS is treated.

Medication is only part of the solution, and in the case of a functional gastrointestinal disorder such as IBS, holistic treatment means addressing all aspects of the disorder, including from the psychological perspective, said Dr Wong.

Psychological interventions that have been used on IBS patients and proven to work through research studies include cognitive behaviour therapy – which helps patients explore how their thinking and emotions are linked – hypnotherapy and a few other modalities, he said.

Stress management, assertiveness training and relaxation exercises, such as progressive muscle relaxation, can also help, said Dr Kwok.

He added that medication, including selective serotonin reuptake inhibitors such as prozac, can help by modifying gut movements and pain sensations, besides reducing anxiety and improving mood.

But not all hospitals have brought the psychologist or psychiatrist on board to see IBS patients.

At Tan Tock Seng Hospital, for example, doctors find that most of their patients respond well to lifestyle modifications and medication.

Dr Robert Lo, a consultant at the department of gastroenterology and hepatology at TTSH, said that once diagnosis of IBS is confirmed, doctors will explain and reassure patients that it is not a life-threatening condition, does not cause damage to the bowel and does not cause cancer.

"Knowledge and awareness are important as they help IBS sufferers to be less anxious, which may help their symptoms," he said.

Dietary advice will be given. Eating a healthy diet with enough fluid and fibre can reduce IBS symptoms. For some people, certain food items, such as alcohol, caffeine, milk products, wheat, citrus fruit and onions, can trigger symptoms.

"IBS sufferers may find it helpful to keep a food diary to identify their food triggers," Dr Lo said.

Exercise is recommended as it can alleviate IBS symptoms. Although the precise cause is not clear, it is thought that exercise relieves stress, which, in turn, reduces IBS symptoms, Dr Lo added.

Only the few who remain symptomatic despite treatment are referred to TTSH psychiatrists.

"We do not see IBS patients routinely with the psychiatrist or psychologist," Dr Lo said.

For Madam Theresa, after she started seeing the psychologist, her physical and mental health has improved.

"I didn't think at first that going for psychological counselling and talking to a psychologist would help me. I also didn't want people to think I was mad. But I am glad that I did because psychotherapy has lessened my IBS pain," she said.

wanching@sph.com.sg

Is IBS getting in the way of your marriage?
Page 14

Is IBS getting in the way of your marriage?

An encounter with the frustrated husband of a patient with irritable bowel syndrome (IBS) led Dr Reuben Wong to wonder about the burden borne by spouses because of the chronic gut condition.

The patient had just left his consultation room when there was a knock on the door. Her husband came inside alone and pleaded with Dr Wong to help his wife get better.

"He told me her symptoms were affecting their marital life negatively, that they no longer had a social life and that he was suffering too," said Dr Wong.

It left a deep impression on the consultant at the department of gastroenterology and hepatology at National University Hospital about the far-reaching impact of IBS.

"When a family member is unwell, we know it affects everyone else in the family too, especially if it's a severe chronic illness," he said.

The phenomenon of partner burden has been described in cancer and dementia, but never explored in gastrointestinal diseases or functional disorders, said Dr Wong.

Not long after, in 2008, Dr Wong left for a postgraduate fellowship at the University of North Carolina at Chapel Hill in the United States.

There, he embarked on a study comparing the partners of 152 IBS patients against the partners of 39 healthy individuals.

The study found that the partners of IBS patients suffered significantly more burden than partners of healthy people.

The patients' partners completed questionnaires including the Zarit Burden Interview

(ZBI), Relationship Satisfaction Scale and questions on sexual relationships.

The ZBI is a previously validated scale which has also been used to measure the burden suffered by the partners or caregivers of people with dementia, cancer and other conditions.

Partners of those with IBS scored 22.1 on the ZBI total score, compared with 11.5 for the control group and 18.5 for partners of cancer patients.

One in three partners also said they believed that IBS interfered in their sex lives either quite frequently or nearly all the time.

The study also showed that the more severe the IBS, the worse the burden reported by the partner or spouse, said Dr Wong.

But if the couple's relationship was stronger, the perceived burden was less.

The study was published in the *Journal Of Clinical Gastroenterology & Hepatology* earlier this year.

Its authors were doctors from the Center for Functional GI and Motility Disorders and department of biostatistics at the University of North Carolina at Chapel Hill and the department of gastroenterology and hepatology of the University Medical Cluster of the National University Health System in Singapore.

Dr Wong, who returned here in 2009, is now hoping to carry his research further.

"In the pipeline is a study to be conducted here, which not only looks at partner burden in Singapore IBS patients, but also evaluates the intervention that will help alleviate this burden," he said.

IBS patients' symptoms are due to an interaction of multiple factors, such as micro-inflammation, stress, previous infections and hypersensitivity of the nerves. These all interact to give the symptoms which are the hallmarks of the condition, said Dr Wong.

facebook

Fans on how they cope with gut problem

I have been suffering from irritable bowel syndrome (IBS) since last year. At first, I thought it was indigestion, as the symptoms included diarrhoea, cramps and bloating. When the symptoms persisted, I saw a doctor and had all sorts of tests, before being told it was IBS.

It hits me on and off. It affects my concentration and quality of life, socially and at work, as it is inconvenient and uncomfortable.

I explain my condition to my colleagues and friends so that they will understand when I am unable to perform my best. I have started paying more attention to my health, exercising and making time to relax, avoiding problem food such as chilli, coffee, milk and cabbage and eating regularly and in moderation.

Chris Hua

Suffering from bowel disorder makes my life miserable and affects my work, especially when I need to stay longer in the toilet. My work gets delayed and I cannot do things properly.

I cope by drinking lots of water and eating apples as it makes my stomach strong. I also rub some oil on my abdominal area.

Rhamhona Cabanting Joseph

I have been experiencing IBS for several years. The symptoms include bloating and alternating periods of diarrhoea and constipation.

Whenever I have diarrhoea, I feel an urgent need to go to the toilet immediately. This has greatly affected my life and I have reduced the frequency of my overseas trips, especially long road trips as I may face difficulty locating toilets in foreign lands.

I have tried several remedies, such as reducing the intake of dairy products and fatty food and having low-fat and high carbohydrate meals. I also keep a food diary so as to keep track of what types of food cause the symptoms.

Diet and lifestyle changes are important in decreasing the frequency and severity of IBS symptoms. Besides taking medication, I also maintain a less stressful lifestyle and practise good relaxation techniques, such as yoga and deep breathing.

Mary Leong

I go crazy whenever I am constipated for a few days. I have tried consuming more water and fruit but it has not helped.

The urge to empty my bowels will strike me at the wrong time in the MRT, bus or car, and you know what happens next...

Lisa Chew

I have always had IBS. I tried taking prune juice but it does not help. I was advised by the doctor to eat spinach and bananas to improve my bowel movements but that has not helped either. I avoid taking any medication as I do not want to rely on it.

However, after reducing stress in my life and including more vegetables and water in my diet, my bowel movement has improved. I still occasionally feel abdominal discomfort. I just let it be.

Sherry Wong

I used to experience frequent bloating, heart burn and flatulence. It caused me to burp uncontrollably. It was very embarrassing for me in the office and the symptoms were uncomfortable and frustrating. It affected my work performance and mood.

Now, I chew my food slowly. Meat is harder to digest so I eat less meat to help my digestive system perform better. When I do eat meat, I make sure that I chew it thoroughly.


I have also changed my diet to include more vegetables and fruit. My IBS symptoms have reduced as a result.

Jeekeng Chew

WINNER

Mary Leong wins a \$50 shopping voucher for the best post. The winner should e-mail her full name, address, e-mail address, identity card number and contact details to shealth@sph.com.sg by next Wednesday. Specify ST MYB Facebook as the subject.

www.facebook.com/STMindYourBody



ST FILE PHOTO